

Exercise Class 1

Exercise 1

Calculate all possible values of the angular momentum L , of its projection on the z-axis L_z , and of the magnetic dipole momentum along z μ_z of an electron when $l = 2$, and plot them graphically.

Exercise 2

A qubit is localized on the Bloch sphere by angles $\theta = 50^\circ$ and $\phi = 10^\circ$. Draw the state on the Bloch sphere and calculate the state vector in the $\{|0\rangle, |1\rangle\}$ basis and the probability of measuring the basis states.

Exercise 3

Consider a qubit $|\psi\rangle = \left(\frac{1}{2} + \frac{i}{2}\right)|0\rangle - \left(\frac{1}{2\sqrt{2}} + i\frac{\sqrt{3}}{2\sqrt{2}}\right)|1\rangle$.

- Locate the state on the Bloch sphere by calculating the corresponding angles θ, ϕ .
- Calculate the global rotation angle δ and the equivalent state $|\psi'\rangle$ with purely real α' coefficient.

Exercise 4

Consider two states $|\psi\rangle, |\psi'\rangle$ differing only by a global phase factor $e^{i\gamma}$. Show that the probability of measuring a state $|s\rangle$ is identical for $|\psi\rangle$ and $|\psi'\rangle$, for any target state $|s\rangle$.

Exercise 5

Consider the Stern-Gerlach experimental setup in Fig. 1, where the input qubit is prepared in state $|\psi\rangle = \frac{1}{\sqrt{2}}|0\rangle - \left(\frac{1+\sqrt{3}}{4} - i\frac{1-\sqrt{3}}{4}\right)|1\rangle$.

- Calculate the measurement probability for states $|x_+\rangle, |x_-\rangle$ and $|y_+\rangle, |y_-\rangle$ after the corresponding experimental setups.
- Suppose now to collimate both the $|x_+\rangle, |x_-\rangle$ output beams into the second SG setup. Calculate the measurement probability for $|y_+\rangle, |y_-\rangle$ after the second SG setup.

